

Your Name

Your Signature

Instructor Name

Problem	Total Points	Score
1	10	
2	10	
3	10	
4	10	
5	10	
6	10	
7	10	
8	10	
9	10	
10	10	
Extra Credit	(5)	
Total	100	

- This test is closed notes and closed book.
- You may **not** use a calculator.
- In order to receive full credit, you must **show your work**. Be wary of doing computations in your head. Instead, write out your computations on the exam paper.
- Raise your hand if you have a question.

1. (10 points) Let  $f(x) = x^2 - \frac{3}{x}$ .

- (a) Using the **definition of the derivative**, find  $f'(x)$ . No credit will be given if a different method is used. [It is recommended you start by writing the definition of the derivative.]

- (b) Write an equation of the line tangent to the graph of  $f(x)$  when  $x = 2$ .

2. (10 points) During a storm, snow is falling on a mountain at a rate of

$$M(t) = t^2 - \frac{t^3}{3}$$

feet per hour for a three hour period starting at time  $t = 0$ .

- (a) Determine the *net change* in the height of snow during the first two hours of the storm. Include units with your answer.

- (b) Determine the height of the snow on the mountain or explain why this is not possible with the present information.

- (c) Observe that  $M(2.5) > 0$  and  $M'(2.5) < 0$ . Explain what these two facts indicate about the snow falling when  $t = 2.5$ .

3. (10 points) The population of ants in a new colony is modeled by the function

$$p(t) = 1000 \left( t + \frac{1}{2} \ln(1 + t^2) \right) + 100,$$

where  $t$  is measured in months.

- (a) Find  $p(0)$  and interpret in the context of the problem.


- (b) Find  $\lim_{t \rightarrow \infty} p(t)$  and interpret in the context of the problem.

- (c) Find  $p'(t)$ .

- (d) Find  $p'(1)$  and interpret in the context of the problem.

- (e) Find  $\lim_{t \rightarrow \infty} p'(t)$  and interpret in the context of the problem.

4. (10 points) Consider the function  $A(x)$  graphed below. Between  $x = 3$  and  $x = 5$ , the graph is of a semicircle of radius 1.



- (a)  $\lim_{x \rightarrow -2} A(x) =$
- (b)  $A(-2) =$
- (c)  $A'(-1) =$
- (d) At what  $x$  values, if any, does  $A'(x)$  not exist?
- (e) Evaluate  $\int_{-1}^2 A(x) dx$ .
- (f) Let  $H(x) = \int_0^x A(s) ds$ . What is the value of  $H(4)$ ?
- (g) For  $H(x)$  from part f., what is the value of  $H'(4)$ .

5. (10 points) Sketch the graph of a function that satisfies **all** of the given conditions. **Label all important items in your graph.**


(a) The domain of  $f$  is  $(-\infty, \infty)$ .

(b)  $f'(x) > 0$  if  $x \neq 2$ .

(c)  $f''(x) > 0$  if  $x < 2$  and  $f''(x) < 0$  if  $x > 2$ .

(d)  $f(2) = 5$

(e)  $\lim_{x \rightarrow \infty} f(x) = 8$  and  $\lim_{x \rightarrow -\infty} f(x) = 0$




6. (10 points) Differentiate the following functions. For parts (a) and (b), it is not necessary to simplify your answers.

(a)  $f(x) = \frac{e^{2x}}{\sqrt{x^2 + 1}} + \arctan(3x)$

(b)  $g(x) = \int_x^2 t \cos(2t^2) dt$

(c) Find  $\frac{dy}{dx}$  by implicit differentiation:  $\ln(xy) - \cos y = ye^x$

7. (10 points) A piece of wire 40 cm long is to be cut to make at most two squares. See the picture below.



- (a) Write the combined area of the two squares as a function of  $x$ . State the domain.
- (b) For what value(s) of  $x$  does the area function have a potential maximum or minimum?
- (c) Where should you cut to minimize the combined area? Maximize the combined area? Justify the classification of the extrema.



8. (10 points)

A particle is moving with acceleration


$$a(t) = t + e^{t/2}.$$

You measure that at time  $t = 0$ , its position is given by  $s(0) = 3$  and its velocity is given by  $v(0) = 8$ .

Determine the position function  $s(t)$ .

9. (10 points)

Consider the function  $g(x) = x\sqrt{4-x^2} - 1$ , part of whose graph is shown below.



- (a) Write down, but do not evaluate, a computation that approximates  $\int_0^2 g(x) dx$  using three **right-hand** rectangles. Draw and shade the rectangles on the graph. (No need to simplify your answer.)

- (b) Determine  $\int_0^2 g(x) dx$  exactly. Show your work, and simplify your answer.

- (c) How could you make the estimate in (a) more accurate?

10. (10 points) Newton's method can be used to find an approximate solution to the equation  $x^2 = 8$ . To apply Newton's method to find these roots, let  $f(x) = x^2 - 8$ .

(a) Use Newton's method with initial approximation  $x_0 = 2$  to find  $x_1$ , a better estimate of a root of the given equation.

(b) Apply one more iteration of Newton's method to find  $x_2$ .

(c) Notice the equation  $x^2 = 8$  has two roots. What value of  $x_0$  would make a good choice to find the **other** root?

**Extra Credit** (up to 5 points):

- (a) Show that the equation  $\cos x - 2x = 4$  has at least one real solution.
- (b) Show that the equation  $\cos x - 2x = 4$  has exactly one real solution. (Hint: what does the derivative tell you about the behavior of the function  $g(x) = \cos(x) - 2x - 4$ ?)