# Fall 2023

Midterm 1

# Math F251X

Name: Solutions

## **Rules:**

- Partial credit will be awarded, but you must show your work.
- You may have a single handwritten  $3'' \times 5''$  notecard, both sides.
- Calculators are not allowed.
- Place a box around your FINAL ANSWER to each question where appropriate.
- Turn off anything that might go beep during the exam.

#### Good luck!

Problem	Possible	Score
1	16	
2	12	
3	10	
4	12	
5	12	
6	8	
7	8	
8	10	
9	12	
Extra Credit	5	
Total	100	

#### 1. (16 points)

The entirety of a function H(x) is shown below. Use the graph of H(x) to answer each question below. If a limit is infinite, indicate that with  $\infty$  or  $-\infty$ . If a value does not exist or is undefined, write **DNE**.



**k**. List the values of x in the domain of H where H(x) is NOT continuous.

 $x = \frac{-2}{3}$  (X=1 is not in the domain)

2

Compute the following limits. If the limit does not exist, write **DNE** and a few words about why it does not exist. If the limit increases without bound, write  $\infty$  or  $-\infty$ .

a. 
$$\lim_{n \to 3} \frac{2x^2 - 5x - 3}{x^2 + x - 12}$$

$$= \lim_{k \to 3} \frac{(2x - 1)(x - 3)}{(x + 4)(x - 3)}$$

$$= \lim_{k \to 3} \frac{(2x - 1)(x - 3)}{(x + 4)(x - 3)}$$

$$= \lim_{k \to 3} \frac{2x - 1}{(x + 4)(x - 3)}$$

$$= \lim_{k \to 3} \frac{2x - 1}{x + 4} = \frac{2(3) - 1}{3 + 4} = \frac{7}{7} = \prod$$
Nope !
Nop

v1

#### 3. (10 points)

Consider the function

$$f(x) = \frac{1}{6-x}.$$

Find f'(5) using the **limit definition of the derivative** and show your work using all appropriate notation. No credit will be awarded for using other methods. Begin by writing down the limit definition of the derivative.

$$f'(5) = \lim_{h \to 0} \frac{f(5+h) - f(5)}{h}$$

$$= \lim_{h \to 0} \frac{1}{h} \left( \frac{i}{6 - (5+h)} - \frac{1}{6 - 5} \right)$$

$$= \lim_{h \to 0} \frac{1}{h} \left( \frac{1}{1 - h} - \frac{1}{1} \right)$$

$$= \lim_{h \to 0} \frac{i}{h} \left( \frac{1 - (1 - h)}{1 - h} \right)$$

$$= \lim_{h \to 0} \frac{i}{h} \left( \frac{1 - (1 - h)}{1 - h} \right)$$

$$= \lim_{h \to 0} \frac{1}{h} \left( \frac{1 - 1 + h}{1 - h} \right)$$

$$= \lim_{h \to 0} \frac{1}{h} \left( \frac{1 - 1}{1 - h} \right)$$

$$= \lim_{h \to 0} \frac{1}{h} \left( \frac{1 - h}{1 - h} \right)$$

$$\begin{array}{c} Chech: \frac{d}{dx}\left(\frac{1}{6-x}\right) = \frac{1}{(6-x)^2} & \text{so at } x = 5 & \text{we get } \frac{1}{(6-5)^2} = 1\\ \dim \frac{1}{h}\left(\frac{1}{6-(k+h)} - \frac{1}{6-x}\right) = \dim \frac{1}{h}\left(\frac{6-x}{(6-(x+h))(6-x)}\right) \\ = \lim_{h \to 0} \frac{1}{h}\left(\frac{6-x-6+x+h}{(6-(x+h))(6-x)}\right) = \lim_{h \to 0} \frac{1}{k}\left(\frac{1}{(6-(x+h))(6-x)}\right) = \frac{1}{(6-x)^2} \\ = \frac{1}{4} & \text{vl} \end{array}$$

#### 4. (12 points)

A police station is located on a straight east-west road. At 9:00 AM, a patrol car leaves the station going east. The function s(t) gives the position of the car, in miles, t hours after 9:00 AM. A positive value for s(t) means that the car is to the east of the station. The graph of s(t) is shown below.



**a**. Determine s(1) and interpret this value in the context of the problem. Your answer should be a sentence and it should include units.

At IDAM, the patrol car is 30 miles east of the station.

**b**. Find the average rate of change between t = 1 and t = 5, and interpret this value in the context of the problem. Your answer should be a sentence and it should include units.

 $\frac{s(5)-s(1)}{5-1} = \frac{-25-30}{4} = \frac{-55}{4}$ Between 10 AM and 2PM, the car was traveling at an average velocity of 55/4 mph west, S(i) = 30S(5) = -25c. Estimate s'(1). Show some work. I need the slope of the tangent dine af(1, s(i)). I will use the points (1.5,40) and (0,10):  $\frac{40-10}{1.5} = \frac{30}{2} = \frac{60}{3}$ - 20

**d**. Explain in simple terms what s'(1) indicates in the context of the problem. Your answer should be a sentence and it should include units.

At the instant of 10 AM, the car is travelling at an instantaneous velocity of 20 miles/how (east) e. In the context of the problem, what happens to the patrol car at t = 4?

The car passes the station at IPM

f. What is s'(2), and what does that mean in the context of the problem?

SI(2)=0, which means that at 11AM the Car stopped (for an instant!) (it was turning around.)

#### 5. (12 points)

Consider the function  $f(x) = \frac{x^2}{x-2}$ .

**a**. Find f'(x). Use whatever method you like. Show your work.

$$f'(x) = \frac{(x-2)(2x) - x^{2}(1)}{(x-2)^{2}} = \frac{2x^{2} - 4x - x^{2}}{(x-2)^{2}} = \frac{x^{2} - 4x}{(x-2)^{2}}$$

**b**. Use your answer from part (a) to determine the slope of the line tangent to f(x) at the point P(1, -1).

$$f'(1) = \frac{1^2 - 4(1)}{(1 - 2)^2} = \frac{1 - 4}{(1 - 2)^2} = \frac{-3}{1} = -3$$

**c**. Write down an equation for the tangent line at the graph of f(x) at the point P(1, -1).

y = -3(x-1)+(-1)

**d**. Clearly draw the tangent line at the graph of f(x) at the point P(1, -1) on the graph below and label it as "tangent".



#### 6. (8 points)

A function g(x) is shown. Sketch the derivative g'(x) on the second set of axes. Indicate any asymptotes the derivative might have using dashed lines, and indicate any points where the derivative is undefined using open circles.



### 7. (8 points)

Consider the function  $f(x) = 2 \sin x - x$ . Determine all *x*-values on the interval  $[0, 2\pi]$  for which f(x) has a **horizontal tangent line**.

$$f'(x) = 2\cos(x) - 1$$

$$f'(x) = 0 \implies 2\cos(x) = 1 \implies \cos(x) = \frac{1}{2}$$

$$\implies x = \frac{\pi}{3} \quad or \quad x = \frac{5\pi}{3}$$

#### 8. (10 points)

Let

$$f(x) = \begin{cases} \frac{3x^2 + x}{x} & x < 0\\ 2 & x = 0\\ \sqrt{x} + e^x & x > 0 \end{cases}$$

**a**. Evaluate  $\lim_{x\to 0^-} f(x)$ . Show supporting work.

$$\lim_{X \to 0^{-}} f(x) = \lim_{X \to 0^{-}} \frac{3x^{2} + x}{x} = \lim_{X \to 0^{-}} 3x + 1 = 1$$

**b.** Evaluate  $\lim_{x\to 0^+} f(x)$ . Show supporting work.

$$\lim_{X \to 0^+} f(x) = \lim_{X \to 0^+} \sqrt{X} + e^X = 0 + 1 = 1$$

**c**. Evaluate f(0).

$$f(0) = 2$$

- d. Based on your answers to parts (a), (b) and (c), check the true statement(s) below:
  - $\Box$  *f* is continuous at x = 0.
  - f has a removable discontinuity at x = 0.
  - $\Box$  *f* has a jump discontinuity at x = 0.
  - $\Box$  *f* has an infinite discontinuity at x = 0.
  - $\Box$  None of the above.

v1

### 9. (12 points)

For each of the following functions, compute the derivative.

## You do not need to simplify your answers.

a. 
$$y = 11x^3 - \frac{4x}{3} + x^2 + x^{5/3}$$
  
 $y' = 11(3x^2) - \frac{4}{3} + 2x + \frac{5}{3} \times \frac{2}{3}$ 

**b.** 
$$a(\theta) = \theta^3 \cos(\theta)$$

$$a'(\theta) = \theta^{3}(-\sin\theta) + \cos\theta(3\theta^{2})$$

c. 
$$f(t) = t\sqrt{t} - \frac{1}{8t^4} + \sqrt{2} = t^{3/2} - \frac{1}{8}t^{-4} + \sqrt{2}$$
  
 $f'(t) = \frac{3}{8}t^{1/2} - \frac{1}{8}(-4t^{-5}) + 0$ 

Extra Credit (5 points) Use the Intermediate Value Theorem to show that the equation  $x^3 = 2^x$  has a solution for some *x*-value. Justify your answer with words (as well as computations).

Consider the function 
$$f(x) = x^3 - 2^x$$
.  
Observe that  
(i)  $f(x)$  is continuous (as the difference  
of continuous functions), and  
(ii) a solution to  $x^3 = 2^x$  is also a  
solution to  $f(x) = 0$ .

Notice  

$$f(a) = 0^3 - a^0 = 0 - 1 = -1 < 0$$
  
 $f(1) = 1^3 - 2^1 = 1 - 2 = -1 < 0$   
 $f(2) = 2^3 - 2^2 = 8 - 4 = 4 > 0$ 

So by the intermediate value theorem, direct f(0) < 0 and f(z) > 0, three must be some value c in (0, 2) g.t. f(c) = 0. This value c is a bolishing that we are looking for!