

Fall 2024

Math F251X

Calculus I: Final Exam

Name: _____

- Section: 9:15 (James Gossell)
 11:45 (Jill Faudree)
 11:45 (Leah Berman)
 Online (James Gossell)

Rules:

- Partial credit will be awarded, but you must **show your work**.
- You may have a single handwritten $3'' \times 5''$ notecard, both sides.
- Calculators are **not allowed**.
- Place a box around your **FINAL ANSWER** to each question where appropriate.
- Turn off anything that might go beep during the exam.
- You have two hours to complete the exam.

Problem	Possible	Score
1	10	
2	6	
3	16	
4	6	
5	8	
6	9	
7	11	
8	9	
9	16	
10	9	
Extra Credit	(5)	
Total	100	

1. (10 points)

Compute the following **integrals**. Give the most general answer, and show your work. Clearly indicate any substitutions you use in such a way that someone else can follow your work.

a. $\int \left(5t^{\frac{2}{7}} - 7t^{-1} + e^{3t-4} + \sin\left(\frac{\pi}{6}\right) \right) dt$


b. $\int x \cos(x^2) + \frac{1}{\sqrt{1-x^2}} dx$

2. (6 points)

Use linearization to estimate $\sqrt{98}$. Show your work, and write your answer as a single decimal or fraction.

3. (16 points)

Consider the graph of the function $f(x)$ shown below, and answer the following questions. You must give the most complete answer; if the value is infinite, write $+\infty$ or $-\infty$.




- a. $\lim_{x \rightarrow 2^-} f(x) = \underline{\hspace{2cm}}$ b. $\lim_{x \rightarrow 6} f(x) = \underline{\hspace{2cm}}$ c. $\lim_{x \rightarrow 2^+} f'(x) = \underline{\hspace{2cm}}$
- d. List all x -values where the derivative $f'(x)$ is not defined. $\underline{\hspace{10cm}}$
- e. Estimate $f'(8) = \underline{\hspace{2cm}}$. **Explain** how you computed your estimate.

- f. List all intervals where $f'(x) > 0$. $\underline{\hspace{10cm}}$
- g. As what x -values does $f(x)$ have...(If none write "none").
 A local maximum? $\underline{\hspace{10cm}}$ A local minimum? $\underline{\hspace{10cm}}$.
- h. The line $y = 0$ is a horizontal asymptote of $f(x)$. Fill in a statement about a limit that corresponds to this fact.

$$\lim_{\boxed{\hspace{1cm}}} \boxed{\hspace{1cm}} = \boxed{\hspace{1cm}}$$
- i. Does f have an(y) inflection point(s)? If so, list it/them, if not write "none."
 Inflection point(s): $x = \underline{\hspace{2cm}}$
- j. On what interval(s) is $f(x)$ concave down? $\underline{\hspace{10cm}}$

4. (6 points)

The graph of a function $g(x)$ is shown below.



Define a new function $A(x) = \int_{-4}^x g(t) dt$.

a. On the interval $(-4, 4)$, does $A(x)$ have a local maximum or a local minimum? If so, give the corresponding x -value and **explain** your answer. If not, explain why not.


b. **Estimate** $A(0) = \underline{\hspace{2cm}}$ and clearly explain what calculation you did to arrive at that estimation. (You may want to draw on the graph as part of your explanation.)

c. Determine $A'(3) = \underline{\hspace{2cm}}$.

5. (8 points)

Below is the graph of the curve $x^4 + y^4 + 2x^2y = 21$.

- a. Find $\frac{dy}{dx}$ for the curve $x^4 + y^4 + 2x^2y = 21$.



- b. Write an equation of the line tangent to the curve $x^4 + y^4 + 2x^2y = 21$ at the point $(1, 2)$.

Tangent line equation: _____

- c. Draw the tangent line on the figure above.

6. (9 points)

The Mars rover drops a rock off of a 61 foot cliff. While in free fall, the rock's **velocity** after t seconds is given by the function $v(t) = -12.2t$ feet per second.

- a. Evaluate the integral $\int_0^2 v(t) dt$ and write a sentence to interpret its meaning in the context of the problem. Include units in your answer.

- b. Write a function $h(t)$ that gives the rock's height above the ground after t seconds.

- c. What is the **acceleration** due to gravity on Mars? Include units in your answer.

7. (11 points)

A box has a square base and an open top. The material for the base costs \$4 per square meter and the material for the sides costs \$1 per square meter. Suppose the width of the base of the box is x meters and its height is y meters.

- a. What is the total cost, C , of the box?

- b. What is the volume, V , of the box?

- c. Suppose you have \$36 to spend on the materials for the box.
 - (i) Write the volume V as a function of one variable and pick a **domain** for this function.

 - (ii) Determine the **dimensions** of the box of **largest** possible volume that fits within your budget.

dimensions (include units): $x =$ _____ $y =$ _____

- (iii) **Justify** that your dimensions give the largest volume, using calculus. As part of your justification, **write the name** of the test you are applying (first derivative test, second derivative test, closed interval/extreme value theorem, some other test).

8. (9 points)

Compute the following **limits**. Show your work clearly. Make sure you use **limit notation** where required; an answer that does not use proper notation will not receive full credit. Use = to show things are equal. If you use L'Hôpital's rule, write $\stackrel{H}{=}$ or $\stackrel{L'H}{=}$ to indicate where you are applying it.

a. $\lim_{h \rightarrow 0} \frac{\ln(e+h) - 1}{h}$

b. $\lim_{\theta \rightarrow \pi} \frac{\sin^2(\theta)}{1 + \cos(\theta)}$

c. $\lim_{x \rightarrow \infty} \frac{-8x^3 + 5x^2}{2x^3 + 3x - 5}$

10. (9 points)


Compute the following **derivatives**. Show your work. You do NOT need to simplify your answer. Your answer should start $f'(x)$, $\frac{df}{dx}$ etc.

a. $f(x) = x^{-4} + \sqrt[4]{2x} - 4^x + 4$

b. $g(x) = \ln\left(\frac{x^5 e^x}{\sqrt{x}}\right)$

c. $h(x) = 5 \sec(2x^{-3})$

Extra Credit (5 points) A portion of the graph of the function $f(x) = x - 5 \arctan(x)$ is shown below.



- Suppose Newton's method is used to find an approximate solution to $f(x) = 0$ from an initial guess of $x_1 = 4$. **Sketch** on the graph how the next approximation x_2 will be found, **labeling** its location on the x -axis.
 - If your starting guess is $x_1 = 4$, **compute** x_2 . Show your work. **You do not need to simplify completely**, but your answer should be in a form where typing it into a calculator would compute a numerical value.
- c. What happens if you try to apply Newton's method with a starting guess of $x_1 = 2$?