

Name: Solutions**Rules:**

You have 90 minutes to complete the exam.

Partial credit will be awarded, but you must show your work.

You may have a single handwritten  $3 \times 5$  notecard.

Calculators are not allowed.

Place a box around your FINAL ANSWER to each question where appropriate.

Turn off anything that might go beep during the exam.

Good luck!

Problem	Possible	Score
1	12	
2	20	
3	10	
4	8	
5	12	
6	15	
7	6	
8	17	
Extra Credit	5	
Total	100	

1. (12 points)

(a) State the definition of  $f'(x)$ , the derivative of the function  $f(x)$ .

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

(b) Find the derivative of  $f(x) = 4 - \sqrt{x}$  using the limit definition of the derivative. No credit will be awarded for using other methods.

$$f'(x) = \lim_{h \rightarrow 0} \frac{(4 - \sqrt{x+h}) - (4 - \sqrt{x})}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\sqrt{x} - \sqrt{x+h}}{h} \cdot \frac{(\sqrt{x} + \sqrt{x+h})}{(\sqrt{x} + \sqrt{x+h})}$$

$$= \lim_{h \rightarrow 0} \frac{x - (x+h)}{h(\sqrt{x} + \sqrt{x+h})} = \lim_{h \rightarrow 0} \frac{-h}{h(\sqrt{x} + \sqrt{x+h})}$$

$$= \lim_{h \rightarrow 0} \frac{-1}{\sqrt{x} + \sqrt{x+h}} = \frac{-1}{2\sqrt{x}}$$

2. (20 points) Evaluate the following limits. Show your work to earn full credit. Be careful to use proper notation.

$$(a) \lim_{x \rightarrow 3} \frac{x^2 - 9}{2x^2 - 5x - 3} = \lim_{x \rightarrow 3} \frac{(x-3)(x+3)}{(x-3)(2x+1)} = \lim_{x \rightarrow 3} \frac{x+3}{2x+1} = \frac{3+3}{2 \cdot 3+1} = \frac{6}{7}$$

$\underbrace{\hspace{10em}}_{\text{form } \frac{0}{0}}$

$$(b) \lim_{x \rightarrow c} \frac{\frac{1}{c} - \frac{1}{x}}{x - c} = \lim_{x \rightarrow c} \frac{\frac{x-c}{cx}}{x-c} = \lim_{x \rightarrow c} \frac{1}{cx} = \frac{1}{c^2}$$

$$(c) \lim_{x \rightarrow 4^-} \frac{\sqrt{x}}{x^2 - 2} = \frac{\sqrt{4}}{4^2 - 2} = \frac{2}{16-2} = \frac{2}{14} = \frac{1}{7}$$

$$(d) \lim_{x \rightarrow -1^-} \frac{8 - 8x}{1 - x^2} = -\infty$$

$\uparrow$  form  $\frac{16}{0}$

As  $x \rightarrow -1^-$  (so  $x$ -values like  $-1.001$ )

$$1 - x^2 \rightarrow 0^- \quad \text{and} \quad 8 - 8x \rightarrow 16$$

So the quotient is  $-\infty$ .

3. (10 points) Let  $f(x) = \begin{cases} \frac{\sqrt{x}+5}{2} & x < 9 \\ 6 & x = 9 \\ 3 + e^{x-9} & 9 < x \end{cases}$ .

(a) Show that  $\lim_{x \rightarrow 9} f(x)$  exists.



$$\lim_{x \rightarrow 9^-} f(x) = \lim_{x \rightarrow 9^-} \frac{\sqrt{x}+5}{2} = \frac{\sqrt{9}+5}{2} = \frac{3+5}{2} = 4$$

$$\lim_{x \rightarrow 9^+} f(x) = \lim_{x \rightarrow 9^+} 3 + e^{x-9} = 3 + e^0 = 3 + 1 = 4$$


(b) Determine if  $f(x)$  is continuous at  $x = 9$  and provide a mathematical justification for your conclusion.

Since the left and right side limits exist and are equal, the two-sided limit exists.

4. (8 points) Use the graph of the function  $g(x)$  (drawn on the left below) to sketch the graph of  $g'(x)$  on the set of axes on the right.



5. (12 points) The function  $g(x) = x^2(x - 1) + 4$  is graphed below.



$$g(x) = x^3 - x^2 + 4$$

$$g'(x) = 3x^2 - 2x$$

$$g'(-1) = 3(-1)^2 - 2(-1) \\ = 3 + 2 = 5$$

$$g(-1) = (-1)^3 - (-1)^2 + 4 \\ = -1 - 1 + 4 \\ = 2$$

- (a) Sketch and label the **tangent** line to the graph of  $g(x)$  when  $x = -1$ .

- (b) Write an equation of the tangent line to  $g(x)$  when  $x = -1$ .

point  $(-1, 2)$       line:  $y - 2 = 5(x + 1)$   
 slope  $m = 5$        $y = 2 + 5(x + 1)$

- (c) Find all  $x$ -values where the graph of  $g(x)$  has a horizontal tangent or explain why none exist.

Find  $x$ -values where  $g'(x) = 3x^2 - 2x = 0$

So  $x(3x - 2) = 0$ .

So  $x = 0$  or  $x = 2/3$

6. (15 points) Find the derivative of each function below. You do not need to simplify your answer.

(a)  $f(x) = 2x^5 + \frac{2}{x^5} + 2\sqrt{5} = 2x^5 + 2x^{-5} + 2\sqrt{5}$

$$f'(x) = 10x^4 - 10x^{-6}$$

(b)  $H(\theta) = \theta^{2/3} \sin(\theta)$

$$H'(\theta) = \frac{2}{3} \theta^{-1/3} \cdot \sin(\theta) + \theta^{2/3} \cos(\theta)$$

(c)  $j(x) = \frac{x}{\pi + \cos(x)}$

$$j'(x) = \frac{(\pi + \cos(x))(1) - x(-\sin(x))}{(\pi + \cos(x))^2} = \frac{\pi + \cos(x) + x \sin(x)}{(\pi + \cos(x))^2}$$

7. (6 points) The function  $V(T)$  models the volume of a fixed mass of gas with respect to temperature where volume  $V$  is measured in milliliters (or  $mL$ ) and temperature  $T$  is measured in kelvins (or  $K$ ).

- (a) Interpret in the context of the problem the meaning of  $V(100) = 60$ . Include units.

At a temperature of 100 kelvins, the volume of the gas is 60 mL.


- (b) Interpret in the context of the problem the meaning of  $V'(100) = 0.4$ . Include units.

At a temperature of 100 kelvins, the volume of gas is increasing at a rate of 0.4 mL per kelvin.

- (c) Using the facts that  $V(100) = 60$  and  $V'(100) = 0.4$ , estimate  $V(110)$ . Include units.

$$V(110) = V(100) + 10 \cdot V'(100) = 60 + 10(0.4) = 64 \text{ K.}$$

8. (17 points) A spring is hanging from the ceiling with a mass attached to it. The mass is oscillating vertically with simple harmonic motion. The function  $h(t) = 6 - \cos(t)$  models the height of the spring above the floor starting at time  $t = 0$  where  $h$  is measured in feet and  $t$  is measured in seconds. **For all problems below, include appropriate units.**



$$h(t) = 6 - \cos(t)$$

$$h(\pi) = 6 - \cos(\pi) = 7$$

$$h(0) = 6 - \cos(0) = 5$$

- (a) Find the average velocity of the mass in the time interval  $[0, \pi]$  seconds.

$$\text{avg vel} = \frac{\Delta h}{\Delta t} = \frac{h(\pi) - h(0)}{\pi - 0} = \frac{7 - 5}{\pi} = \frac{2}{\pi} \text{ ft/s}$$

- (b) Find the equations for the velocity and acceleration of the mass.

$$h'(t) = \sin(t) \text{ ft/s}$$

$$h''(t) = \cos(t) \text{ ft/s}^2$$

- (c) Find the instantaneous velocity of the mass at  $\pi/4$  seconds.

$$h'\left(\frac{\pi}{4}\right) = \sin\left(\frac{\pi}{4}\right) = \frac{\sqrt{2}}{2} \text{ ft/s}$$

- (d) At  $t = \pi/4$ , is the mass going up or going down? Justify your answer.

going up

- (e) Determine all times  $t > 0$  at which the mass changes direction?

$$\text{Check for } v'(t) = 0.$$

$$\text{Solve } \sin(t) = 0$$

$$\text{So } t = 0, \pi, 2\pi, 3\pi, \dots$$

**Extra Credit:** (5 points) Use the Intermediate Value Theorem to demonstrate that the function  $f(x) = x^3 - 3x - 19$  has a root (or zero) in the interval  $[1, 5]$ . A complete answer requires some calculations **and** complete sentences justifying your conclusion.

Since  $f(x)$  is a polynomial, it is continuous and the Intermediate Value Theorem applies.

Observe that  $f(1) = 1^3 - 3(1) - 19 < 0$  and

$$f(5) = 5^3 - 3 \cdot 5 - 19 > 0.$$

Since  $f(1) < 0$  and  $f(5) > 0$ , the Intermediate Value Theorem implies that  $f(x) = 0$  for some  $x$ -value between 1 and 5.