

LECTURE: 3-8 EXPONENTIAL GROWTH AND DECAY

In many natural phenomena, a quantity grows or decays at a rate proportional to their size. Suppose $y = f(t)$ is the number of individuals in a population at time t . Given an unlimited environment, adequate nutrition and immunity to disease it is reasonable to assume that the rate of growth is proportional to the population. That is,

$$f'(t) = \frac{dy}{dt} = \frac{k y}{1}$$

\leftarrow constant of proportionality
 \leftarrow population
 \uparrow rate of growth

Example 1: Show that the equation $y = Ce^{kt}$ is a solution to the differential equation $\frac{dy}{dt} = ky$.

(a) Explain, in words, what it means for $y = Ce^{kt}$ to be a solution of the given differential equation.

When you "check" this in $dy/dt = ky$ it will "work".

(b) Show that $y = Ce^{kt}$ is a solution to the differential equation $\frac{dy}{dt} = ky$.

$$\frac{dy}{dt} = C \cdot e^{kt} \cdot k$$

$$\frac{dy}{dt} = k \cdot \underbrace{Ce^{kt}}$$

$$\frac{dy}{dt} = ky !$$

~~~~~

**Theorem:** The only solutions of the differential equation  $dy/dt = ky$  are exponential functions of the form  $y(t) = Ce^{kt}$  where  $C = y(0)$

- Explain why  $C = y(0)$ .

$y(0) = Ce^{k(0)}$  This means that  $C$  is the initial population at time  $t=0$ .  
 $y(0) = C$  ;

- What does the constant  $k$  mean in this equation? What does the sign of  $k$  tell you about the growth of your population?

$k$  is the rate of growth of the population;

If  $k > 0$  the population is growing.

If  $k < 0$  the population is decreasing.

**Example 1:** A bacteria culture initially contains 10 cells and grows at a rate proportional to its size. After an hour the population has increased to 400.

- (a) Find an expression for the number of bacteria after  $t$  hours.  $f(0) = 10$ ,  $f(1) = 400$

$$f(t) = Ce^{kt}$$

$$f(0) = Ce^0$$

$$10 = C$$

$$\text{so } f(t) = 10e^{kt}$$

$$f(1) = 400$$

$$400 = 10e^{k \cdot 1}$$

$$40 = e^k$$

$$k = \ln 40$$

Thus,

$$f(t) = 10e^{(\ln 40)t}$$

$$= 10 \cdot 40^t$$

- (b) Find the number of bacteria after 3 hours.

$$f(3) = 10 \cdot 40^3$$

$$= \boxed{640,000 \text{ bacteria}}$$

- (c) Find the rate of growth after 3 hours.

$$f'(x) = 10e^{\ln 40 x} \cdot \ln 40$$

$$f'(3) = 10 \cdot \ln 40 \cdot 40^3$$

$$= \boxed{23,608,828,506 \text{ bacteria/hour}}$$

- (d) When will the population reach 1,000?

$$1000 = 10e^{\ln 40 t}$$

$$100 = e^{\ln 40 t}$$

$$\ln 100 = \ln 40 t$$

$$t = \frac{\ln 100}{\ln 40}$$

$$\approx \boxed{1.248 \text{ hours}}$$

**Example 2:** Let  $y = Ce^{kt}$  be the number of flies at time  $t$ , where  $t$  is measured in days. Suppose there are 100 flies after the second day and 400 flies after the fourth day. Assuming the growth rate is proportional to the population size find a model for this population's growth. When will the population of flies be 10,000?

$(2, 100)$  and  $(4, 400)$

$$100 = Ce^{2k} \Rightarrow C = \frac{100}{e^{2k}}$$

$$400 = Ce^{4k}$$

$$400 = \frac{100}{e^{2k}} e^{4k}$$

$$400 = 100 e^{2k} \quad C = \frac{100}{4}$$

$$4 = e^{2k}$$

$$C = 25$$

$$\ln 4 = 2k$$

$$k = \frac{1}{2} \ln 4$$

$$y = 25 e^{(\frac{1}{2} \ln 4) \cdot t}$$

$$= 25 (e^{\ln 4^{1/2}})^t$$

$$y = 25 (2)^t$$

$$10000 = 25 \cdot 2^t$$

$$4000 = 2^t$$

$$\ln(4000) = \ln 2^t$$

$$\ln(4000) = t \ln 2$$

$$t = \frac{\ln 4000}{\ln 2} \approx 11.966 \text{ days}$$

**Example 3:** The half-life of cesium-137 is 30 years. The 1986 explosion at Chernobyl sent about 1000 kg of radioactive cesium-137 into the atmosphere.

(a) Find the mass that remains after  $t$  years.

$$m(t) = 1000 \left(\frac{1}{2}\right)^{t/30}$$

$$m(t) = Ce^{kt}$$

$$m(t) = 1000 e^{kt}$$

know  $t=30, m(30)=500$

$$500 = 1000 e^{30k}$$

$$\frac{1}{2} = e^{30k}$$

$$\ln\left(\frac{1}{2}\right) = 30k$$

$$k = \frac{1}{30} \ln\left(\frac{1}{2}\right)$$

$$m(t) = 1000 e^{\frac{1}{30} \ln\left(\frac{1}{2}\right) t} = 1000 \left(e^{\ln\left(\frac{1}{2}\right)^{1/30}}\right)^t$$

$$= 1000 \left(\left(\frac{1}{2}\right)^{1/30}\right)^t$$

$$= 1000 \left(\frac{1}{2}\right)^{t/30}$$

(b) If even 100 kg remains in Chernobyl's atmosphere, the area is considered unsafe for human habitation. Determine when Chernobyl will be safe.

$$100 = 1000 \left(\frac{1}{2}\right)^{t/30}$$

$$10 = \left(\frac{1}{2}\right)^{t/30}$$

$$\ln(10) = \ln\left(\frac{1}{2}\right)^{t/30}$$

$$\ln(10) = \frac{t}{30} \ln\left(\frac{1}{2}\right)$$

$$t = \frac{30 \ln(10)}{\ln\left(\frac{1}{2}\right)} \approx 99.658 \approx 100$$

→ 100 years after 1986,  
so  $\boxed{2086}$

**Example 4:** A sample of radioactive tritium-3 decayed to 95% of its original amount after a year.

(a) What is the half-life of tritium-3?

$$y = C \left(\frac{1}{2}\right)^{(t/h)}$$

$$t=1, y=(0.95)C$$

$$\text{so, } 0.95C = C(0.5)^{1/h}$$

$$0.95 = (0.5)^{1/h}$$

$$\ln(0.95) = \ln(0.5)^{1/h}$$

$$\ln(0.95) = \frac{1}{h} \ln(0.5)$$

$$h \ln(0.95) = \ln(0.5)$$

$$h = \ln(0.5) / \ln(0.95)$$

$$h \approx 13.513$$

(b) How long would it take the sample to decay to 10% of its original amount?

$$0.1C = C(0.5)^{t/13.513}$$

$$0.1 = (0.5)^{t/13.513}$$

$$\ln(0.1) = \frac{t}{13.513} \ln(0.5)$$

$$t = \frac{13.513 \ln(0.1)}{\ln(0.5)}$$

$$t \approx 44.891 \text{ years}$$

**Example 5:** Scientists can determine the age of ancient objects by the method of *radiocarbon dating*. The bombardment of the upper atmosphere by cosmic rays converts nitrogen to a radioactive isotope of carbon,  $^{14}\text{C}$ , with a half life of about 5730 years. Vegetation absorbs carbon dioxide through the atmosphere and animal life assimilates  $^{14}\text{C}$  through food chains. When a plant or animal dies, it stops replacing its carbon and the amount of  $^{14}\text{C}$  begins to decrease through radioactive decay. Therefore, the level of radioactivity must also decay exponentially.

A parchment fragment was discovered that had about 74% as much  $^{14}\text{C}$  radioactivity as does the plant material on earth today. Estimate the age of the parchment.

$$f(t) = C \left(\frac{1}{2}\right)^{t/5730}$$

Q: when is  $f(t) = 0.74C$ ?

$$0.74C = C(0.5)^{t/5730}$$

$$0.74 = (0.5)^{t/5730}$$

$$\ln(0.74) = \ln(0.5)^{t/5730}$$

$$\ln(0.74) = \frac{t}{5730} \ln(0.5)$$

$$t = \frac{5730 \ln(0.74)}{\ln(0.5)}$$

$$t \approx 2489.13 \text{ years}$$

old @ discovery

### Newton's Law of Cooling

The rate of cooling (or warming) of an object is proportional to the temperature difference between the object and its surroundings:

$$\frac{dT}{dt} = k(T - T_s)$$

temp  $\rightarrow$   $T$        $\leftarrow$  temp of surroundings  $T_s$

**Example 6:** When a cold drink is taken from a refrigerator, its temperature is 40° F. After 25 minutes in a 70° F room its temperature has increased to 52° F.

(a) What is the temperature of the drink after 50 minutes?

not quite  $\frac{dy}{dt} = ky$ ; close  $\rightarrow$  let  $y = T - T_s$ , this works

Have  $y = T - 70$  here;  $y(0) = 40 - 70 = -30$

Thus  $y(t) = -30 e^{kt}$ , also after  $t=25$  min drink is 52° F

so  $y(25) = 52 - 70 = -18$  and ...

$$-18 = -30 e^{25k}$$

$$\frac{9}{15} = e^{25k}$$

$$25k = \ln\left(\frac{9}{15}\right)$$

$$k = \ln\left(\frac{9}{15}\right)/25$$

$$y(t) = -30 e^{(\frac{1}{25} \ln(9/15))t}$$

$$T - 70 = -30 e^{\frac{1}{25} \ln(9/15)t}$$

$$T(t) = 70 - 30 e^{\frac{1}{25} \ln(9/15)t}$$

$$T(50) = 59.2^\circ\text{F}$$

$$t \approx 87.689 \text{ min}$$

(b) When will its temperature reach 65° F?

$$65 = 70 - 30 e^{\frac{1}{25} \ln(9/15)t}$$

$$-5 = -30 e^{\frac{1}{25} \ln(9/15)t}$$

$$\frac{1}{6} = e^{\frac{1}{25} \ln(9/15)t}$$

$$\ln(1/6) = \frac{1}{25} \ln(9/15)t$$

$$\frac{25 \ln(1/6)}{\ln(9/15)} = t$$

(c) What happens to the temperature of the drink as  $t \rightarrow \infty$ ? Is this expected?

$$\lim_{t \rightarrow \infty} T(t) = \lim_{t \rightarrow \infty} (70 - 30 e^{-0.6385t})$$

$$= \lim_{t \rightarrow \infty} (70 - \frac{30}{e^{0.6385t}})$$

$= 70^\circ\text{F}$   $\leftarrow$  as  $t \rightarrow \infty$ ,  $T(t) \rightarrow 70^\circ\text{F}$  or the ambient room temp.