

1. State, formally, the definition of the derivative of a function  $f(x)$  at  $x = a$ .

$$\lim_{a \rightarrow x} \frac{f(x) - f(a)}{x - a} \quad \text{or} \quad \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

2. Let  $f(x) = 5x^2 - 3x$ .

1. Use the definition to find the derivative of  $f(x)$ .

$$\begin{aligned} f'(x) &= \lim_{a \rightarrow x} \frac{f(x) - f(a)}{x - a} = \lim_{a \rightarrow x} \frac{5x^2 - 3x - (5a^2 - 3a)}{x - a} \\ &= \lim_{a \rightarrow x} \frac{5(x^2 - a^2) - 3(x - a)}{x - a} \\ &= \lim_{a \rightarrow x} \frac{5(x - a)(x + a) - 3(x - a)}{x - a} \\ &= \lim_{a \rightarrow x} 5(x + a) - 3 = 5(a + x) - 3 \end{aligned}$$

2. Find the slope of the tangent line to  $f(x)$  when  $x = -3$ .

$$f'(-3) = -33$$

$$= \boxed{10a - 3}$$

3. Write the equation of the line tangent to  $f(x)$  when  $x = -3$ .

$$\begin{aligned} y &= f(-3) + f'(-3)(x - (-3)) \\ &= 54 - 33(x + 3) \end{aligned}$$

3. Suppose  $N$  represents the number of people in the United States who travel by car to another state for a vacation this Memorial Day weekend when the average price of gasoline is  $p$  dollars per gallon.

1. What are the units of  $dN/dp$ ?

people/dollar



2. In the context of the problem, interpret  $\frac{dN}{dp}$ .

This is the rate at which the number of travellers changes as the price of gas increases.

3. Would you expect  $dN/dp$  to be positive or negative? Explain your answer.

Negative. The number of travelers should decrease as the price of gas goes up.

4. The graph of  $f(x)$  is sketched below. On a separate set of axes, give a rough sketch  $f'(x)$ .



5. Find the domain of each function.

1.  $f(x) = \sqrt{x^2 - x - 6}$

2.  $g(t) = \ln(t + 6)$

Need  $x^2 - x - 6 \geq 0$   
 $(x+2)(x-3) \geq 0$

So  $x \leq -2$  or  $x \geq 3$

$(-\infty, -2] \cup [3, \infty)$

$t + 6 > 0$

$t > -6$

$(-6, \infty)$

6. State the definition of "The function  $f(x)$  is continuous at  $x = a$ ".

$$\lim_{x \rightarrow a} f(x) = f(a)$$

7. Suppose

$$f(x) = \begin{cases} -\frac{2}{x} & x < 2 \\ \frac{x}{x-3} & x \geq 2 \end{cases}$$

Is  $f(x)$  continuous at  $x = 0$ ? At  $x = 2$ ? Justify your answers using the definition of continuity.

At 0? No. The function isn't defined there.

At  $x=2$ ? No.  $\lim_{x \rightarrow 2^-} -\frac{2}{x} = -1$ ,  $\lim_{x \rightarrow 2^+} \frac{x}{x-3} = \frac{2}{-1} = -2$ .

Since  $-1 \neq -2$ ,  $\lim_{x \rightarrow 2} f(x)$  does not exist, much less equal  $f(2)$ .

8. Find the limit or show that it does not exist. Make sure you are writing your mathematics correctly and clearly.


$$1. \lim_{x \rightarrow \infty} \frac{10^x - 1}{3 - 10^x} = \lim_{x \rightarrow \infty} \frac{1 - 10^{-x}}{3 \cdot 10^{-x} - 1} = \frac{1 - 0}{0 - 1} = -1$$

$$\begin{aligned} 2. \lim_{x \rightarrow \infty} \frac{\sqrt[3]{8x^3 + 1}}{2 - 5x} &= \lim_{x \rightarrow \infty} \frac{x^3 \sqrt[3]{8 + 1/x^3}}{2 - 5x} \\ &= \lim_{x \rightarrow \infty} \frac{\sqrt[3]{8 - 1/x^3}}{2/x - 5} \\ &= \frac{\sqrt[3]{8}}{-5} = \frac{-2}{5} \end{aligned}$$

9. Write the formula for a function with vertical asymptotes at  $x = -1$  and  $x = 3$  and a horizontal asymptote at  $y = 4/3$ .

$$f(x) = \frac{1}{(x+1)(x-3)} + \frac{4}{3}$$

10. Sketch the graph of the function from problem 7.



Note:

$$\begin{aligned} \frac{x}{x-3} &= \frac{x-3}{x-3} + \frac{3}{x-3} \\ &= 1 + \frac{3}{x-3} \end{aligned}$$

11. Solve for  $x$ .

1.  $e^{x-3} + 2 = 6$

$$e^{x-3} = 4$$

$$x-3 = \ln(4)$$

$$x = 3 + \ln(4)$$

2.  $\ln(x+5) - 3 = 7$

$$\ln(x+5) = 10$$

$$x+5 = e^{10}$$

$$x = e^{10} - 5$$

3.  $\ln x + \ln(x-1) = 0$

$$\ln(x(x-1)) = 0$$

$$x(x-1) = e^0 = 1$$

$$x^2 - x - 1 = 0 \quad x = \frac{1 \pm \sqrt{1+4}}{2}$$

4.  $\cos(8x) = 0$

$$8x = \frac{\pi}{2} + k\pi$$

$$x = \frac{\pi}{16} + k\frac{\pi}{8}$$

$$k \in \mathbb{Z}$$

$$= \frac{1 \pm \sqrt{5}}{2}$$


But  $\frac{1-\sqrt{5}}{2}$

is not in the domain.

12.

1. What does the Intermediate Value Theorem say? You may want to include a picture with your explanation.

If  $f(x)$  is continuous on  $[a, b]$  and  $c$  is a number between  $f(a)$  and  $f(b)$  then there is  $x$  in  $[a, b]$  with  $f(x) = c$ :




2. Use the Intermediate Value Theorem to show  $\ln x = x - 5$  has a solution. (Hint: Show there is a solution in the interval  $[1, e^5]$ .)

Let  $f(x) = \ln(x) - x + 5$ . Notice  $f(x)$  is continuous on  $(0, \infty)$  and so also on  $[1, e^5]$ . Moreover,  
 $f(1) = 0 - 1 - 5 = -6 < 0$  and  
 $f(e^5) = \ln(e^5) + e^5 - 5 = e^5 > 0$ .

13. So there is  $x$  in  $[1, e^5]$  with  $f(x) = 0$ .

1. What does the Squeeze Theorem say? You may want to include a picture with your explanation.

If  $g(x) \leq f(x) \leq h(x)$  near  $x=a$ , but maybe not at  $x=a$ , and if  $\lim_{x \rightarrow a} g(x) = \lim_{x \rightarrow a} h(x) = L$ , then  $\lim_{x \rightarrow a} f(x) = L$  also.



14. Use the Squeeze Theorem to show  $\lim_{x \rightarrow \infty} \frac{\cos(2x)}{3x^2} = 0$ .

Since  $-1 \leq \cos(2x) \leq 1$ ,  $\frac{-1}{3x^2} \leq \frac{\cos(2x)}{3x^2} \leq \frac{1}{3x^2}$ .

Since  $\lim_{x \rightarrow \infty} \frac{-1}{3x^2} = \lim_{x \rightarrow \infty} \frac{1}{3x^2} = 0$ ,  $\lim_{x \rightarrow \infty} \frac{\cos(2x)}{3x^2} = 0$ .

15. Sketch each of the functions below. Label all  $x$ - and  $y$ -intercepts and asymptotes. State, in interval notation, the domain and range of each function next to its graph.

1.  $y = 6 - x^4$

4.  $y = \tan^{-1} x$

7.  $y = -2/(x + 3)$

2.  $y = \sin(2x)$

5.  $y = e^{x-1} + 2$

8.  $y = \sqrt{x+5}$

3.  $y = \tan x$

6.  $y = \ln x$

