

## 4-3

**Mean Value Theorem.** If  $f$  is a continuous function on an interval  $[a, b]$  that has a derivative at every point in  $(a, b)$ , then there is a point  $c$  in  $(a, b)$  where

$$f'(c) = \frac{f(b) - f(a)}{b - a}.$$

1. Suppose  $f$  is a continuous function on  $[a, b]$  that has a derivative at every point of  $(a, b)$ . Suppose also that  $f(b) \leq f(a)$ . What can you conclude from the Mean Value Theorem?

2. Suppose  $f$  is a continuous function on  $[a, b]$  that has a derivative at every point of  $(a, b)$ , and that  $f'(x) > 0$  for each  $x$  in  $(a, b)$ . Thinking about your answer to problem 1, what can you conclude about  $f(a)$  and  $f(b)$ ?

3. A function is said to be **increasing** on an interval  $(a, b)$  if whenever  $x$  and  $z$  are in the interval and  $x < z$ , then  $f(x) < f(z)$ . It is **decreasing** if whenever  $x$  and  $z$  are in the interval and  $x < z$ , then  $f(x) > f(z)$ . Sketch an example of a function that is increasing on  $(1, 3)$  and decreasing on  $(3, 5)$ .

### Increasing/Decreasing Test

Your answer to problem 2 implies the first item below; the second is justified by a similar argument.

- If  $f'(x) > 0$  on an interval  $(a, b)$  then  $f$  is increasing on the interval.
- If  $f'(x) < 0$  on an interval  $(a, b)$  then  $f$  is decreasing on the interval.

4. Use the increasing/decreasing test to find intervals where

$$f(x) = \frac{2}{3}x^3 + x^2 - 12x + 7$$

is increasing and intervals where it is decreasing.

5. Find the critical points of the function  $f(x) = \frac{2}{3}x^3 + x^2 - 12x + 7$  from the previous problem. There should be two,  $c_1$  and  $c_2$  with  $c_1 < c_2$ . Just pay attention to  $c_1$ .

- Just to the left of  $c_1$  is the function increasing or decreasing?
- Just to the right of  $c_1$  is the function increasing or decreasing?
- Now decide intuitively, based on these two observations, if  $f$  has a local min, local max, or neither at  $c_1$ .

6. Repeat the previous exercise for the other critical point  $c_2$ .

You have just sketched the argument that justifies the following:

**First Derivative Test**



Suppose  $f$  is a function with a derivative on  $(a, b)$ , and if  $c$  is a point in the interval with  $f'(c) = 0$ .

- If  $f'(x) > 0$  for  $x$  just to the left of  $c$  and  $f'(x) < 0$  for  $x$  just to the right of  $c$ , then  $f$  has a \_\_\_\_\_ at  $c$ .
- If  $f'(x) < 0$  for  $x$  just to the left of  $c$  and  $f'(x) > 0$  for  $x$  just to the right of  $c$ , then  $f$  has a \_\_\_\_\_ at  $c$ .

7. The function  $f(x) = xe^x$  has exactly one critical point. Find it, and then use the First Derivative Test to determine if a local minimum or local maximum occurs there.

8. Consider the function  $f(x) = \frac{2}{3}x^3 + x^2 - 12x + 7$ . Find intervals such that the **derivative** of  $f(x)$  is increasing or decreasing.

9. Earlier you computed that  $f'(-3) = 0$ . Is  $f'$  increasing near  $x = -3$  or decreasing near  $x = -3$ ? Which of the following two scenarios must we have?



You have just sketched out justification for the following.

**Second Derivative Test**

Suppose  $f$  is a function with a continuous second derivative on  $(a, b)$ , and that  $c$  is a point in the interval with  $f'(c) = 0$ .

- If  $f''(c) > 0$  then  $f$  has a \_\_\_\_\_ at  $c$ .

- If  $f''(c) < 0$  then  $f$  has a \_\_\_\_\_ at  $c$ .

10. Use the Second Derivative Test to determine if  $f(x) = xe^x$  has a local min/max at its only critical point.

11. Consider the function  $f(x) = x^3$ . Verify that  $f'(0) = 0$ . Then decide what the Second Derivative Test has to say, if anything, about whether a local min/max occurs at  $x = 0$ .

12. Decide what the First Derivative Test has to say, if anything, about whether a local min/max occurs at  $x = 0$  for  $f(x) = x^3$ .