

Name: \_\_\_\_\_

\_\_\_\_\_ / 25

There are 25 points possible on this quiz. No aids (book, calculator, etc.) are permitted. **Show all work for full credit.**

1. [10 points] Let  $P(0, 1)$  be a point on the graph of  $f(x) = \sqrt{x+1}$ .

a. Find the **slope of the secant line** passing through  $P$  and the point  $Q(3, f(3))$ .


b. The table below lists the slope ( $m_{sec}$ ) of the secant line passing through the point  $P$  and the point  $Q(x, f(x))$  for several values of  $x$ .

$x$	-1	-0.1	-0.01	0.01	0.1	1
$f(x)$	0	0.9487	0.9950	1.0050	1.0488	1.4142
$m_{sec}$	1.0	0.5132	0.5013	0.4988	0.4881	0.4142

Use the information in the table to estimate the **slope of the tangent line** to  $f(x)$  at the point  $P(0, 1)$ .

c. Use the slope from part (b) above to write an **equation of the tangent line** at point  $P(0, 1)$ .

d.



Left is a sketch of the graph of  $f(x) = \sqrt{x+1}$ .

**Sketch and label the tangent line** to the graph at the point  $P(0, 1)$ .

**Sketch and label the secant line** between  $P(0, 1)$  and  $Q(3, f(3))$ .

2. [5 points] A professional cyclist is riding along a straight road. For the first minute, the distance in feet that the cyclist has traveled after  $t$  seconds is given by the function  $p(t) = \frac{1}{2}t^2 + t$ . Find the **average velocity** of the cyclist between  $t = 2$  and  $t = 4$  seconds. Include units with your answer.

3. [8 points] Evaluate the expressions below. Assume all angles are measured in radians.


a.  $\cos(\pi/4) =$

b.  $\sin(7\pi/6) =$

c.  $\tan(\pi/3) =$

d.  $\sin(-\pi/2) =$

4. [2 points] Use the right triangle below, with side lengths 12, 5, and 13, to evaluate the expressions.



a.  $\cot(\theta) =$

b.  $\sec(\theta) =$